

The Adolescent Knee and Risk for Osteoarthritis — An Opportunity or Responsibility for Sport Medicine Physicians?

Sharon Bout-Tabaku¹ and Thomas M. Best²

¹Nationwide Children's Hospital and The Ohio State University College of Medicine, Columbus, OH; ²OSU Sports Medicine Center, The Ohio State University College of Medicine, Columbus, OH

INTRODUCTION

Knee osteoarthritis (OA) is a degenerative joint disease that affects articular cartilage, bone, and synovium. It most commonly occurs in adults over the age of 50 yr, affecting women more than men. Knee OA, unfortunately, can occur in younger adults who are in their 30s, often through sport-related joint injury (19,32). Exceptionally, degenerative joint disease can occur in children or adolescents with a syndromic disease such as Stickler syndrome, Kniest syndrome, or spondyloepiphyseal dysplasia (6). Knee OA risk factors and disease mechanisms include mechanical, systemic, and genetic causes. A recent meta-analysis including 49 prospective studies showed that body mass index (BMI), previous knee injury, female gender, intensive physical activity, certain occupational activities, and increased bone mineral density are risk factors for onset of knee OA (4). Obesity and joint injury from sports participation in young adults are known risk factors that lead to osteoarthritis onset in adults, and both risk factors are on the rise among adolescents today (11,29,31). Thus, knee OA is a chronic disease that may evolve over years and involve multiple risk factors that may be present in young adults and possibly even in adolescence. Accordingly, sports medicine practitioners need to counsel young patients and their families about the risks for knee OA.

Osteoarthritis is the fifth leading cause of disability in older Americans today (2). With the increasing prevalence of musculoskeletal complaints and sports injuries in children at younger ages, are we to expect an increase in the prevalence of knee OA, and will it occur at an earlier age? Not surpris-

ingly, there is an impetus to understand the natural history and significance of adolescent risk factors on incidental knee OA in adulthood. Understanding this relationship may lead to identifying at-risk adolescents and lead to evidence-based interventions targeted at preventive and disease modifying treatments to decrease the burden of knee OA. Here we present the evidence on adolescent sports-related trauma and incident knee OA, the association of sports on joint injury in adolescence, biomechanical risk factors for incidental knee OA, and promising treatment approaches.

SPORTS INJURIES AND KNEE OA

The evidence to date is limited to young adults and clearly links knee OA to anterior cruciate ligament (ACL), meniscal, and articular cartilage injuries. Among young adults, knee injury was shown to lead to knee OA with a cumulative incidence of 13.9% at age 65 compared to 6.0% in those without knee injury (11). Subsequent prospective studies found radiographic osteoarthritis developing within 12 to 15 yr in female and male soccer players with ACL and meniscal damage (20,32). In long-term follow up of young athletes who had meniscus surgery, more than 50% developed knee OA, pain, and functional impairment (22).

The data on children and adolescents and whether they develop osteoarthritis in childhood due to sports injuries is sparse. One study in children, aged 4 to 17 yr, who had internal derangements of the knee found chondral lesions to be prevalent, and more so in skeletally immature children than in older mature children. However, there was no follow up on the long-term resolution of these early lesions and whether they progress to osteoarthritis (27).

SPORTS-RELATED INJURIES IN THE ADOLESCENT

Epidemiologic data demonstrate that sport injuries cause acute morbidity among adolescent athletes and at alarmingly

Address for correspondence: Sharon Bout-Tabaku, M.D., Nationwide Children's Hospital and The Ohio State University College of Medicine, 700 Children's Drive, Columbus, OH 43205 (E-mail: Sharon.bout-tabaku@nationwidechildrens.org).

increasing rates in the ankle and the knee (7). Most commonly, football, wrestling, basketball, and soccer are the cause of injuries that occur at the knee, ankle, and ligaments (9). Compounding the already-high adolescent injury rate is the added burden of obesity in children, which leads to a threefold risk of musculoskeletal injuries (23,35). In an environment where sports participation is greater, at a higher intensity level, and occurring at a younger age, the risk of mechanical knee injuries in adolescence is great and sets the stage for knee OA.

KNEE BIOMECHANICS AND OA

The biomechanical causes of osteoarthritis arise from traumatic ligament and cartilage injuries, leading to abnormal loading of the joint that causes alteration of shear and compression forces. Thus the disease process ensues with disruption and fibrillation of cartilage, change in the cartilage matrix, and involvement of underlying bone (33). Articular cartilage damage also can occur as a result of repetitive joint loading (8) that causes blunt impact on the tissue and subsequent matrix degradation and chondrocyte death (5,16). Animal models demonstrate that, even at impact energies not great enough to sustain articular fracture, progressive chondrocyte death occurs specifically in areas of thinner cartilage (30). Thus increased loading forces across the joint and articular cartilage lead to remodeling by way of an accelerated healing that occurs in response to tissue injury known as the “regional acceleratory phenomenon.” The chondral response curve further explains how cartilage responds to increasing load 1) initially accelerated longitudinal growth of the limb which reaches a peak and then 2) at larger loads decreased or cessation of growth (10). Thus for child and adolescent athletes whose bones and cartilage are developing, added stress and loading forces may alter the normal joint development and serve as an additional risk factor for knee OA.

TREATMENT

As known risk factors for knee OA (such as sports-related injury and obesity) are becoming more prevalent in children and adolescents, sports medicine personnel have an opportunity to modify their approach to injury management and counseling on health lifestyles. Treatment of these injuries can be addressed through both primary and secondary interventions. To date, primary prevention strategies have focused on altering lower extremity biomechanics. A meta-analysis revealed that multiintervention training programs were effective in reducing the risk of lower limb injuries by 39%, acute knee injuries by 54%, and ankle sprains by 50% (13). Another systematic review concluded that using insoles, external joint supports, and training programs were effective (1).

Secondary treatment of the adolescent athlete traditionally has been limited to surgical intervention. Treatment of ACL and meniscal injuries has focused on tissue repair and reconstruction, with much attention paid to the impact of these

procedures on epiphyseal damage in skeletally mature versus immature children but little attention paid to long-term outcomes (22). The protective role of ACL reconstruction for osteoarthritis onset was accepted by many; however, systematic review of randomized control trials (RCT) showed a lack of evidence to support that surgical management is better (18,24). Adjunctive surgical intervention such as functional tissue engineering for ACL injuries with growth factors, gene therapy, cell therapy, and extracellular matrix scaffolds have been used in human subjects and lab models (12). For cartilage injuries, marrow stimulating therapies, microfracture of subchondral bone to release pleuripotent cells, growth factors to regenerate tissue, osteochondral autograft, autologous chondrocyte implantation, and osteochondral allografts all have been suggested for prevention of osteoarthritis (21,25).

Chondroprotective pharmacological therapies, commonly used in adults with knee OA, may offer promise in adolescents sustaining significant joint injuries. The early use of cyclooxygenase inhibitors, nonionic surfactant, or derivatives of glucosamine following joint injury has shown to reduce chondrocyte death (14,15,26). Recent Osteoarthritis Research Society International (OARSI) recommendations for osteoarthritis demonstrated that intraarticular hyaluron had the highest effect sizes for improvement in pain, function, and stiffness (34). Chondroitin sulfate had no significant effect sizes for pain, but one RCT demonstrated a statistically significant decline in the rate of reduction of joint space narrowing over 2 yr in the treatment arm (17). The effect of intraarticular corticosteroid injection showed some effect over the first year with regard to pain but did not have any effect on function and stiffness. Glucosamine sulfate had a small effect size for pain but was even smaller when only higher-quality trials were included, while its effect on structural modification remains controversial (34).

The evidence for chondroprotective therapies in sports injuries is less robust, with fewer studies, small numbers, and short follow up. In one randomized study (treatment or no treatment) with 58 d of follow up, there was improvement with pain and mobility in the intraarticular sodium hyaluronate treatment arm for acute knee injuries with mild to moderate ligament tears and instability (3). More recently, a 2-yr follow-up study of athletes with acute ankle sprain showed that hyaluronic injections compared with placebo were more effective in reducing pain and reinjury (28). Long-term follow up to assess the impact of chondroprotection for sports injuries on the risk of incidental knee OA has not been studied.

In conclusion, we know that athletic knee injuries can predispose to knee OA. Further, more children are playing sports at a higher intensity level and are incurring more injuries. Whether these injuries put the child at higher risk for developing knee OA at an earlier age remains an important and unresolved issue. However, the treating physician should be cognizant of the potential for later onset degenerative disease in the younger athlete and particularly should be concerned with children who have multiple risk factors for adult onset osteoarthritis, such as female gender and obesity. Finally, while primary prevention is important, newer chondroprotective approaches may be of clinical use and require more research in the younger athlete.

References

1. Aaltonen S, Karjalainen H, Heinonen A, *et al.* Prevention of sports injuries: systematic review of randomized controlled trials. *Arch. Intern. Med.* 2007; 167(15):1585–92.
2. American Association of Orthopaedic Surg. *Burden of Musculoskeletal Diseases in the United States: Prevalence, Societal and Economic Cost.* Rosemont (IL): American Association of Orthopaedic Surg; 2008.
3. Axe MJ, Shields CL Jr. Potential applications of hyaluronans in orthopaedics: degenerative joint disease, surgical recovery, trauma and sports injuries. *Sports Med.* 2005; 35(10):853–64.
4. Blagojevic M, Jinks C, Jeffery A, Jordan KP. Risk factors for onset of osteoarthritis of the knee in older adults: a systematic review and meta-analysis. *Osteoarthritis Cartilage.* 2010; 18(1):24–33.
5. Borrelli J Jr, Tinsley K, Ricci WM, *et al.* Induction of chondrocyte apoptosis following impact load. *J. Orthopaedic Trauma.* 2003; 17(9):635–41.
6. Cassidy J, Petty RE, Laxer RM, Linsley CB. *Textbook of Pediatric Rheumatology.* Philadelphia: Elsevier; 2005.
7. Cuff S, Loud K, O'Riordan MA. Overuse injuries in high school athletes. *Clin. Ped.* 2010; 49(8):731–6.
8. Curl WW, Krome J, Gordon ES, *et al.* Cartilage injuries: a review of 31,516 knee arthroscopies. *Arthroscopy.* 1997; 13(4):456–60.
9. Darrow CJ, Collins CL, Yard EE, Comstock RD. Epidemiology of severe injuries among United States high school athletes: 2005–2007. *Am. J. Sports Med.* 2009; 37(9):1798–805.
10. Frost HM. Biomechanical control of knee alignment: some insights from a new paradigm. *Clin. Orthop. Relat. Res.* 1997; (335):335–42.
11. Gelber AC, Hochberg MC, Mead LA, *et al.* Joint injury in young adults and risk for subsequent knee and hip osteoarthritis. *Ann. Intern. Med.* 2000; 133(5):321–8.
12. Hsu SL, Liang R, Woo SL. Functional tissue engineering of ligament healing. *Sports Med. Arthrosc. Rehabil. Ther. Technol.* 2010; 2:12.
13. Hubscher M, Zech A, Pfeifer K, *et al.* Neuromuscular training for sports injury prevention: a systematic review. *Med. Sci. Sports Exerc.* 2010; 42(3):413–21.
14. Huser CA, Davies ME. Effect of a glucosamine derivative on impact-induced chondrocyte apoptosis in vitro. A preliminary report. *Osteoarthritis Cartilage.* 2008; 16(1):125–8.
15. Jeffrey JE, Aspden RM. Cyclooxygenase inhibition lowers prostaglandin E2 release from articular cartilage and reduces apoptosis but not proteoglycan degradation following an impact load in vitro. *Arthritis Res. Ther.* 2007; 9(6):R129.
16. Jeffrey JE, Gregory DW, Aspden RM. Matrix damage and chondrocyte viability following a single impact load on articular cartilage. *Arch. Biochem. Biophys.* 1995; 322(1):87–96.
17. Kahan A, Uebelhart D, De Vathaire F, *et al.* Long-term effects of chondroitins 4 and 6 sulfate on knee osteoarthritis: the study on osteoarthritis progression prevention, a two-year, randomized, double-blind, placebo-controlled trial. *Arthritis Rheum.* 2009; 60(2):524–33.
18. Linko E, Harilainen A, Malmivaara A, Seitsalo S. Surgical versus conservative interventions for anterior cruciate ligament ruptures in adults. *Cochrane Database Syst. Rev.* 2005; (2):CD001356.
19. Lohmander LS, Englund PM, Dahl LL, Roos EM. The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. *Am. J. Sports Med.* 2007; 35(10):1756–69.
20. Lohmander LS, Ostenberg A, Englund M, Roos H. High prevalence of knee osteoarthritis, pain, and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis Rheum.* 2004; 50(10):3145–52.
21. Macmull S, Skinner JA, Bentley G, *et al.* Treating articular cartilage injuries of the knee in young people. *BMJ.* 2010; (340):c998.
22. Maffulli N, Longo UG, Gougoulas N, *et al.* Long-term health outcomes of youth sports injuries. *Br. J. Sports Med.* 2010; 44(1):21–5.
23. McHugh MP. Oversized young athletes: a weighty concern. *Br. J. Sports Med.* 2010; 44(1):45–9.
24. Meuffels DE, Favejee MM, Vissers MM, *et al.* Ten year follow-up study comparing conservative versus operative treatment of anterior cruciate ligament ruptures. A matched-pair analysis of high level athletes. *Br. J. Sports Med.* 2009; 43(5):347–51.
25. Minas T, Gomoll AH, Solhpour S, *et al.* Autologous chondrocyte implantation for joint preservation in patients with early osteoarthritis. *Clin. Orthopaed. Relat. Res.* 2010; 468(1):147–57.
26. Natoli RM, Athanasiou KA. P188 reduces cell death and IGF-I reduces GAG release following single-impact loading of articular cartilage. *J. Biomechan. Eng.* 2008; 130(4):041012.
27. Oeppen RS, Connolly SA, Bencardino JT, Jaramillo D. Acute injury of the articular cartilage and subchondral bone: a common but unrecognized lesion in the immature knee. *AJR.* 2004; 182(1):111–7.
28. Petrella MJ, Coglianò A, Petrella RJ. Original research: long-term efficacy and safety of periarticular hyaluronic acid in acute ankle sprain. *Phys. Sportsmed.* 2009; 37(1):64–70.
29. Roos EM. Joint injury causes knee osteoarthritis in young adults. *Curr. Opin. Rheumatol.* 2005; 17(2):195–200.
30. Szczodry M, Coyle CH, Kramer SJ, *et al.* Progressive chondrocyte death after impact injury indicates a need for chondroprotective therapy. *Am. J. Sports Med.* 2009; 37(12):2318–22.
31. Thelin N, Holmberg S, Thelin A. Knee injuries account for the sports-related increased risk of knee osteoarthritis. *Scand. J. Med. Sci. Sports.* 2006; 16(5):329–33.
32. Von Porat A, Roos EM, Roos H. High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes. *Ann. Rheum. Dis.* 2004; 63(3):269–73.
33. Wolf BR, Amendola A. Impact of osteoarthritis on sports careers. *Clin. Sports Med.* 2005; 24(1):187–98.
34. Zhang W, Nuki G, Moskowitz RW, *et al.* OARSI recommendations for the management of hip and knee osteoarthritis: part III: changes in evidence following systematic cumulative update of research published through January 2009. *Osteoarthritis Cartilage.* 2010; 18(4):476–99.
35. Zonfrillo MR, Seiden JA, House EM, *et al.* The association of overweight and ankle injuries in children. *Ambul. Pediatr.* 2008; 8(1):66–9.